

# **Applied math 2<sup>nd</sup> Sec.**

## **Final revision**

The position vector of a moving body is given by  $\vec{r} = (3t + 2)\vec{i} + (4t - 1)\vec{j}$ , then the magnitude of its displacement till  $t = 2$  sec. equals = ..... length unit.

(a) 9

(b) 5

(c) 10

(d) 8

A particle moves such that its position vector  $\vec{r} = (t + 1)\vec{i} + (t - 2)\vec{j}$ , then the magnitude of displacement between two moments  $t = 2$  to  $t = 4$  equals ..... length units.

(a)  $2\sqrt{2}$

(b) 8

(c)  $2\sqrt{10}$

(d) 6

A particle move in a straight line from a fixed point (O) such that its position vector  $\vec{r}$  is given by the relation  $\vec{r} = (t^2 + 3t + 5) \vec{n}$  where  $\vec{n}$  is a unit vector parallel to the straight line. The average velocity after 3 seconds from the beginning of motion is .....  $\vec{n}$

(a) 20

(b)  $\frac{20}{3}$ 

(c) 6

(d) 18

The positions of a moving particle at two instants 3 seconds and 8 seconds was at A (7 , 2) and B (4 , 6) respectively

, then the average velocity of the particle = .....

(a)  $3\hat{i} - 4\hat{j}$

(b)  $-3\hat{i} + 4\hat{j}$

(c)  $\frac{3}{5}\hat{i} - \frac{4}{5}\hat{j}$

(d)  $\frac{-3}{5}\hat{i} + \frac{4}{5}\hat{j}$

A cyclist moved 6 km. to west , then 8 km. in direction of  $60^\circ$  north of west , then magnitude of covered displacement equals ..... km.

(a) 14

(b)  $2\sqrt{37}$

(c)  $2\sqrt{13}$

(d) 2

A cyclist covered 60 km. towards West , then he moved 90 km. towards East , the velocity in two cases was 12 km./h , then average velocity vector is .....

- (a) 12 km./h. West.    (b) 12 km./h. East.    (c) 2.4 km./h. East.    (d) 12.5 km./h. East.

Two cars A and B moves on the same straight road in opposite directions with speeds 125 km./hr. , 75 km./hr. respectively , then the speed of car B relative to the car A = .....

(a) 50

(b) – 50

(c) 200

(d) 75



A motorcycle moves with speed 40 km./hr. in direction of a fixed unit vector  $\hat{c}$ , its rider watches a car, it seems to him that it moves in the opposite direction with speed 105 km./hr., then the velocity of the car is .....

(a)  $155 \hat{c}$

(b)  $-65 \hat{c}$

(c)  $65 \hat{c}$

(d)  $-155 \hat{c}$

A moving radar car to monitor the velocity on the desert road moves with constant velocity 40 km./hr. This car observes the movement of a truck coming in the opposite direction. It seems like it is moving with velocity 120 km./hr. , then the actual velocity for the truck = ..... km./hr.

(a) 160

(b) 80

(c) 120

(d) 40

Two trains A and B , the length of each is 100 m. they are moving in opposite directions with velocities 15 m./sec. and 25 m./sec. , then time taken to cross each other = ..... sec.

(a) 4

(b) 5

(c) 6

(d) 8

A particle moves in a straight line with uniform retardation of magnitude  $3 \text{ m./sec}^2$  to become at rest after 19 seconds. , then the magnitude of the initial velocity = ..... m./sec.

(a) 16

(b) 54

(c) 60

(d) 57

A particle started its motion with velocity  $20 \text{ cm./sec.}$  and uniform acceleration  $8 \text{ cm./sec}^2$  in the same direction as the initial velocity , then the distance covered in the fifth second only = ..... cm.

(a) 200

(b) 144

(c) 100

(d) 56

A particle moves from rest in a straight line with uniform acceleration , it covered 9 cm. at the 5<sup>th</sup> second only , so acceleration = .....  $\text{cm}^2/\text{s}$ .

(a) 2

(b) 3

(c) 4

(d) 6

A body moves in a fixed direction with an initial speed and uniform acceleration. The body covers 20 m. in the third sec. then covers 60 m. in the fifth and sixth seconds , then its initial speed equals ..... m./sec.

(a) 4

(b) 10

(c) 20

(d) 30

If a body fell from a height 19.6 m. above a sandy ground to embed in it a distance 14 cm. till it rests , then the acceleration of the motion of the body inside the sand = ..... m./sec<sup>2</sup>.

(a) – 1372

(b) – 9.8

(c) 19.6

(d) 1732



A body moves with initial velocity ( $v_0$ ) , acceleration ( $a$ ) and final velocity ( $v$ )  
 ,  $v - v_0 = 8 \text{ cm./sec.}$  ,  $v + v_0 = 25 \text{ cm./sec.}$  , then  $\sqrt{4as} = \dots\dots\dots$

- (a)  $10 \text{ cm./sec}^2$       (b)  $20 \text{ cm./sec}^2$       (c)  $20 \text{ cm./sec.}$       (d)  $50 \text{ cm./sec.}$

A body is projected vertically upward with speed 42 m./sec. , then its maximum height the body reach equals ..... m.

(a) 65

(b) 98

(c) 84

(d) 90

A body is projected vertically up from a point on the surface on the ground to return to it after 10 sec. from the instant of projection , the initial velocity = ..... m./sec.

(a) 9.8

(b) 4.9

(c) 98

(d) 49

A body started its motion with velocity 7 m./sec. and with uniform acceleration  $2 \text{ m./sec}^2$ , it covered a distance 30 metres, then the acceleration stopped to move afterwards with uniform velocity a distance 52 metres, then the total time of motion = ..... second.

(a) 3

(b) 4

(c) 7

(d) 14

**In the opposite figure :**


A cyclist moves from (O) in a straight line then :

**First :** Magnitude of the average velocity during the whole journey = ..... m./sec.

- (a) 2                      (b) 4  
(c) 14                    (d) 16

**Second :** The average speed during the whole journey = ..... m./sec.

- (a) 2                      (b) 4  
(c) 14                    (d) 16



From the top of a tower its height 20 metres a particle is projected vertically upward with initial velocity 7 m./sec. , then its speed at the moment of reaching ground surface = ..... m./sec.

(a)  $7\sqrt{7}$

(b) 7

(c) 21

(d)  $7\sqrt{5}$

From the top of a tower , a body is projected vertically upward with initial velocity 9.8 m./sec. , it reached the ground surface after 12 seconds , then the height of the tower = ..... m.

(a) 490

(b) 588

(c) 498

(d) 534

A body is projected vertically upward from the top of a building 32.4 m. high with a velocity of 24 m./sec. , then the elapsed time till it reaches the ground surface = ..... sec.

(a) 4

(b) 8

(c) 5

(d) 6



A body is projected horizontally against direction of wind with initial speed 15 cm./sec. to move with uniform deceleration  $5 \text{ cm./sec}^2$ , then the time elapsed till the body returned to its projected point is .....

(a) 2

(b) 3

(c) 4

(d) 6

A particle moves with initial velocity ( $v_o$ ) cm./sec. , and its final velocity ( $v$ ) cm./sec.

, with acceleration ( $a = 0$ ) , then  $\frac{v^2 + v_o^2}{vv_o} = \dots\dots\dots$

(a)  $2 v_o$

(b)  $\frac{1}{2} v_o$

(c)  $\frac{1}{2}$

(d) 2

A body moves from rest with uniform acceleration for 20 sec. If it covers distance (K) m. in the first 10 seconds and (M) m. in the next 10 seconds , then  $M = \dots\dots\dots$

(a) k

(b) 2 k

(c) 3 k

(d) 4 k

**In the opposite figure :**

$\overline{AB}$  is the hour hand in a clock , if its length is 7 cm.

, then the magnitude of displacement done by the point B

when the hour hand moves from 1 o'clock to 4 o'clock = ..... cm.



- (a)  $\frac{7}{2} \pi$       (b)  $14 \pi$       (c)  $7\sqrt{2}$       (d) 38.5

A body moved from rest with uniform acceleration  $3 \text{ m./sec}^2$ , then the covered distance during  $4^{\text{th}}$ ,  $5^{\text{th}}$  and  $6^{\text{th}}$  seconds = ..... m.

- (a)  $13 \frac{1}{2}$                       (b) 15                      (c) 45                      (d)  $40 \frac{1}{2}$

If a particle moves such that  $\frac{\text{the magnitude of the displacement}}{\text{the covered distance}} = x$ , then .....

(a)  $x = 1$

(b)  $x > 1$

(c)  $x \in [0, 1]$

(d)  $-1 < x < 1$

A satellite of mass 4000 kg. rotates around the earth in an elliptic orbit at a height 440 km. from the ground surface. if the mass of the earth is  $6 \times 10^{24}$  kg. and its radius length =  $6.36 \times 10^6$  metres. , then the gravitational force between the earth and the satellite  $\simeq$  ..... newton. ( $G = 6.67 \times 10^{-11}$  n.m<sup>2</sup>/kg<sup>2</sup>)

- (a) 34418                      (b)  $2.35 \times 10^{11}$                       (c) 34619                      (d) 39575

Two spheres , the mass of the first is  $m$  kg. and the mass of the second is  $\frac{1}{2}m$  kg.

The distance between their centres is 10 m. The universal gravitational constant is  $6.67 \times 10^{-11}$  newton.  $\text{m}^2/\text{kg}^2$  , then the gravitational force = ..... newton.

- (a)  $6.67 \times 10^{-13}$       (b)  $3.335 \times 10^{-13}$       (c)  $3.335 \times 10^{13}$       (d)  $3.533 \times 10^{-13}$



The mass of a planet is three times the mass of the earth and its diameter length equals three times the diameter length of the earth , then the ratio between the gravity acceleration on the surface of this planet and the gravity acceleration on the surface of the earth = .....

(a) 1 : 9

(b) 1 : 3

(c) 9 : 1

(d) 3 : 1

Best wishes  
Mr. Michael Gamil